arXiv:cond-mat/0305264v1 [cond-mat.soft] 12 May 2003

A possible new phase of antagonistic nematogens in a disorienting field

T. G. Sokolovska[1,2],* M. E. Cates[1], and R. O. Sokolovskii[1,2]
[1]*School of Physics, JCMB King's Buildings, University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*
[2]*Institute for Condensed Matter Physics, 1 Svientsitskii, Lviv 79011, Ukraine*

(Dated: February 2, 2008)

A simple model is proposed for nematogenic molecules that favor perpendicular orientations as well as parallel ones. (Charged rods, for example, show this antagonistic tendency.) When a small disorienting field is applied along z, a low density phase N_- of nematic order parameter $S_z < 0$ coexists with a dense biaxial nematic N_b. (At zero field, N_- becomes isotropic and N_b uniaxial.) But at stronger fields, a new phase N_{+4}, invariant under $\pi/2$ rotations around the field axis, appears in between N_- and N_b. Prospects for finding the N_{+4} phase experimentally are briefly discussed.

PACS numbers: 64.70.Md, 61.30.Cz, 61.30.Gd

Practical applications of nematics are based on their strong responses to weak external influences. Therefore, the phase behavior of nematics in external fields of different symmetries is a subject of many investigations (see [1, 2] and references therein). It was shown that external fields can lead to major changes in phase transition behavior. In nematics with positive anisotropy, an external magnetic or electric field is 'orienting': it favors positive nematic order parameter S. In strong orienting fields the first order phase transition between the 'isotropic' phase (which for nonzero field acquires finite $S > 0$) and the nematic phase N is suppressed above a critical temperature. In systems with negative molecular anisotropy, similar fields have instead a disorienting effect, favoring $S < 0$. The isotropic phase then acquires uniaxial planar orientation; with decreasing temperature or increasing density, this phase N_- has a phase transition into a biaxial phase N_b (which connects smoothly to N as the field is removed). A critical field is expected, above which the N_-/N_b phase transition becomes second order [1].

Previous theoretical investigations of external field effects (in cases where the interaction potential was explicitly specified [3]) were carried out [2] for molecular potentials proportional to the second Legendre polynomial of the relative molecular orientation, $P_2(\cos\omega_{12})$, which is the simplest dependence favouring parallel alignment of nematic particles. In practice, though, real molecules have interactions described by more complex orientational dependences than this. For example, uniaxial nematogenic particles may display disposition towards mutually perpendicular orientations as well the parallel one. The molecular interaction then has minima at both $\omega_{12} = 0$ and $\omega_{12} = \pi/2$. (Generically these are of unequal depths.) An example of this antagonistic tendency is for the case of charged rods. It was shown that the Debye-Huckel expression for the electrostatic repulsion between two effective line charges at mutual angle ω_{12} is proportional to $|\sin\omega_{12}|^{-1}$. As a consequence, two charged rods tend to rotate to a perpendicular configuration; this interaction is antagonistic to the usual (steric) Onsager interaction, favoring parallel alignment, which is also present. Such an electrostatic twisting results in significant quantitative changes of the nematic phase diagram in zero field (see Ref. [4] and references therein). Some nonconvex particles, dumbbells for example, should have a certain tendency to mutual twisting too.

The simplest angular dependence of a potential describing molecules with a tendency to orient perpendicularly as well as parallel to each other is the fourth order Legendre polynomial

$$P_4(\cos\omega_{12}) = [35\cos(4\omega_{12}) + 20\cos(2\omega_{12}) + 9]/64 \quad (1)$$

In general, this P_4 interaction should be taken in addition to, not in place of, the customary $P_2(\cos\omega_{12})$ dependence. However, absence of a P_2 term does not imply higher molecular symmetry than when it is present; a pure P_4 interaction already has unequal well depths for perpendicular and parallel interactions.

The task of this paper is to consider an 'antagonistic' nematic in the presence of an external disorienting field. The origin of the disorienting field may vary: electric or magnetic fields, orienting flow in side-chain polymers (with a side group trend to be perpendicular to the backbone), surface influences and so on. The only important factor is that the particles want to orient perpendicularly to the field direction. Note that besides being ubiquitous in display applications, which involve both surface and electric fields [5], the disorienting effect in a sample could be strongly enhanced in some cases by addition of impurities [6].

Our model comprises a system of uniaxial molecules in a disorienting field of strength W, where the potential of the molecular interaction with the field is

$$v(1) = v(\omega_1) = W(3\cos^2\theta_1 - 1)/2, \quad W > 0, \quad (2)$$

$\omega_i = (\theta_i, \varphi_i)$ being the orientation of molecule i. The above formula assumes that the field is directed along the z axis and states that molecular orientations along the field direction are energetically unfavorable. The pair potential is taken to be a sum of the hard sphere potential

---

*Electronic address: tata@icmp.lviv.ua

for spheres of diameter σ, and of an anisotropic part

$$v(1,2) = \sum_{n=2,4} v_n(R_{12}) P_n(\cos\omega_{12}), \tag{3}$$

where ω_{12} is the angle between the axes of molecules 1 and 2; R_{12} is the distance between their centers; and

$$v_n(R_{12}) = -A_n (z_n\sigma)^2 \frac{\exp(-z_n R_{12})}{R_{12}/\sigma}. \tag{4}$$

In the limit $z_n\sigma \to 0$, this model belongs to a family of the so-called Kac potentials, for which the mean field approach is accurate, and the free energy obeys:

$$\begin{aligned} \beta F/N &= F_0 + \int f(\omega)[\ln f(\omega) - 1] d\omega \\ &+ \beta W \int f(\omega) P_2(\cos\theta) d\omega \\ &- \sum_{n=2,4} 12\beta A_n \eta \int f(\omega_1) P_n(\cos\omega_{12}) f(\omega_2) d\omega_1 d\omega_2, \end{aligned} \tag{5}$$

where $\eta = \pi(N/V)\sigma^3/6$, $f(\omega)$ is a single particle distribution function, and for the free energy of the reference hard sphere system we use the "quasiexact" Carnahan-Starling expression, $F_0 = \ln\eta + (4\eta - 3\eta^2)/(1-\eta)^2 + \text{const.}$

To make the mathematics more compact, we give details only for the particular case $A_2 = 0$. (However, adding a $P_2(\cos\omega_{12})$ term is straightforward.) Using a spherical harmonic expansion for $P_n(\cos\omega_{12})$ [7] and noting that the model under consideration is non-polar, one can get the explicit expression for $\Delta F \equiv \beta F/N - F_0$:

$$\begin{aligned} \Delta F &= \int f(\omega)[\ln f(\omega) - 1] d\omega + \beta W \langle P_2(\cos\theta)\rangle_\omega \\ &- \beta A_4 \eta \frac{12}{9} [\langle Y_{40}(\omega)\rangle_\omega^2 + 2\langle Y_{42}^c(\omega)\rangle_\omega^2 + 2\langle Y_{44}^c(\omega)\rangle_\omega^2], \end{aligned} \tag{6}$$

where $\langle\cdots\rangle_\omega = \int f(\omega)(\cdots)d\omega$ denotes orientational averages within a self-consistent one-particle distribution function. The superscript 'c' means that the real part of the spherical harmonic Y_{ij} is taken; (6) is written assuming that the zx-plane is a symmetry plane. This overcomes a continuous degeneracy of the free energy connected with the fact that a spontaneous ordering can rotate without any energy cost in the xy-plane.

Minimizing (6) with respect to $f(\omega)$ we obtain the self-consistent single particle distribution function (with C a normalization constant)

$$\begin{aligned} f(\omega) &= C \exp\{-\beta W \langle P_2(\cos\theta)\rangle_\omega \\ &+ \frac{24}{9}\beta A_4 \eta [\langle Y_{40}(\omega)\rangle_\omega Y_{40}(\omega) \\ &+ 2\langle Y_{42}^c(\omega)\rangle_\omega Y_{42}^c(\omega) + 2\langle Y_{44}^c(\omega)\rangle_\omega Y_{44}^c(\omega)]\}. \end{aligned} \tag{7}$$

We calculated the phase diagrams of this system using, where appropriate, the (standard) common-tangent construction to find coexisting states of different density. The symmetries of all the resulting phases can be defined using three order parameters:

$$\begin{aligned} S_z &= \langle P_2(\cos\theta)\rangle_\omega, & (8) \\ S_x &= \left\langle \sin^2\theta \cos(2\varphi)\right\rangle_\omega, & (9) \\ S_{xy} &= \left\langle \sin^2\theta \cos(4\varphi)\right\rangle_\omega. & (10) \end{aligned}$$

Here S_z and S_x describe alignment of molecules along the corresponding axes, whereas S_{xy} describes ordering along the y axis as well as along the x axis; when $S_x = 0$, S_{xy} describes ordering with fourfold symmetry.

In Fig. 1 the phase diagrams are presented for different values of the ratio W/A_4 describing the external field strength in units of the molecular interaction. (We set $A_2 = 0$ as explained above.) At zero field ($W/A_4 \to -0$) the system undergoes a first order phase transition from an isotropic to a uniaxial nematic N; this phase has $S_z > 0$, $S_x = S_{xy} = 0$. Small disorienting fields ($W/A_4 = 0.5$) transform the system symmetry, so that the isotropic phase at low densities is replaced by a uniaxial planar phase N_- (this has $S_z < 0$, $S_x = S_{xy} = 0$). At intermediate densities (between the bold solid lines in the figure) this coexists with a highly ordered biaxial nematic phase N_b. The biaxial has $S_z < 0$, $S_x > 0$, $S_{xy} > 0$, and becomes a stable single phase at high densities. At higher fields the orientational transition becomes second order at high temperatures, joining onto the first order transition (which is accompanied by the miscibility gap) at the tricritical point. The wide miscibility gap at low temperatures is connected with the appearance of an effective attraction between oriented molecules, which causes the gas liquid phase transition. Molecular potentials of typical thermotropic nematics are characterized by strong isotropic attractions, in addition to anisotropic interactions. In this case the condensation could arise in a non-ordered phase. However, some lyotropic systems display a wide miscibility gap between phases of different symmetries. So far, the picture is qualitatively just as found previously for a pure $P_2(\cos\omega_{12})$ interaction [1]. At low disorienting field strengths, then, the antagonistic tendency of our molecules cannot manifest itself and the behavior is that of a conventional nematic system.

However, at strong enough fields (e.g., $W/A_4 = 2$) the situation differs strongly from the conventional case: two consecutive orientational transitions now take place. The first one is always of second order and meets the binodal at a critical end point. The resulting phase is invariant under rotations by $\pi/2$ around the field direction; it has $S_z < 0$, $S_x = 0$, and $S_{xy} > 0$. In accordance with the general classification [8], this new phase should be denoted as N_{+4}. The second orientational phase transition is from N_{+4} into N_b; this changes from first to second order at the tricritical point as before. The new N_{+4} phase is (like N_b) a result of spontaneous ordering, and is characterized by long-range correlations of Goldstone type. Note that the molecular potential under investigation does not possess the symmetry of the N_{+4} phase (even for the particular case studied, of $A_2 = 0$). In



FIG. 1: The phase diagram of the system ($A_2 = 0$) in (a) the temperature-density ($T^* = k_\mathrm{B}T/A_4$, $\eta = \pi\rho\sigma^3/6$) and (b) pressure-temperature coordinates ($P^* = (\pi\sigma^3/6)(P/A_4)$) for different values of the disorienting field (W/A_4). N_{+4} is the phase with $S_{xy} \neq 0$, $S_x = 0$, N_b denotes the biaxial nematic phase. Dashed lines correspond to the the second order phase transition into N_{+4} phase, thin solid lines likewise into N_b; thick lines denote the first order transition. Circles are tricritical points, diamonds denote critical end points.

TABLE I: Coordinates of the tricritical point ($A_2 = 0$) at different values W/A_4 of the disorienting field. (Notation as in Fig. 1.)

W/A_4	T^*	η	P^*
1	2.02	0.62	39.1
2	1.70	0.48	8.59
5	1.30	0.32	1.52
50	1.52	0.29	1.40

some respects the situation is similar to that found by computer simulations, when a cubatic state appears in a system of hard truncated spheres [10]. The N_{+4} phase is not cubatic but might be called 'square-atic', since it has fourfold rotational order in the transverse plane only.

From the calculated phase diagrams (especially in $T-\eta$ coordinates) one can see a nonmonotonic efect of the external field on the N_{+4} region. The widest N_{+4} region arises at intermediate field values. We attribute this fact to the disorienting field's ability to restrict the part of orientational phase space in which particles mainly reside. Indeed, in the limit of an infinite field, the particle orientations are confined in the plane perpendicular to the field. In this case the system can be described by the plane rotator model [11]; this has the same phase diagram (independent of n) for all pair potentials proportional to $\cos(n\omega_{12})$. In contrast, with an unrestricted orientational phase space (that is, at zero field), the temperature at which spontaneous ordering occurs decreases with increasing n. Disorienting fields (pushing the system towards the plane rotator state) promote the ordering with $S_{xy} \neq 0$ more strongly than that with $S_x \neq 0$, so that the N_{+4} phase grows leaving behind competing ordered phases such as N_b. However, in strong enough fields the susceptibility of the order parameter S_{xy} to the field decreases, whereas the N_b ordering is not saturated. As a result, in strong enough fields the N_b phase begins to displace the N_{+4} region.

This field effect on the N_{+4} region is closely linked with the tricritical behavior (Table I). The nonmonotonic field effect on where the tricritical point lies can be explained by the existence of two competing tendencies [9]. First, the external field furthers the orientational ordering of particles and, therefore, causes a more effective attraction between particles. This favors the phase separation and raises the 'binodal' in temperature. The second tendency takes place if the susceptibility of the low-density phase (N_{+4}) to the field is larger than that of the coexisting dense (N_b) phase. This decreases the energy gain of the phase separation. The second tendency is very strong for our system. Only at very strong fields ($W/A_4 = 50$) do the susceptibilities of the coexisting phases become comparable, allowing the first tendency to win at last.

To study the robustness of the predicted N_{+4} phase with respect to changes of the intermolecular potential we calculated phase diagrams for $A_2/A_4 = 0.1$; 0.2; 0.3 at $W/A_4 = 5$; 10; 50; ∞ in a density interval sensible for liquids ($\eta < 0.6$). In Fig. 2 the phase diagram for $A_2 = 0.1A_4$ is presented (setting $W/A_4 = 5$). With this potential, the parallel orientation of molecules is more energetically favorable than in the case $A_2 = 0$. As a consequence, the N_b transition temperature increases, whereas the temperature of the N_{+4} transition is not affected significantly. The N_{+4} region is reduced in size, but the



FIG. 2: The phase diagram of the model (3) with $A_2 = 0.1A_4$ in the temperature-density and pressure-temperature coordinates for a moderate disorienting field ($W/A_4 = 5$). (Notation as in Fig. 1.) For comparison, the location of the N_{+4} phase for the case $A_2 = 0$ is shaded.

phase remains stable. The phase transition N_{+4}/N_b is first order now (where previously it was second order, for this value of W/A_4) which should lead to a jump in the elastic constants, for example. For all the investigated cases with $A_2 > 0$ the phase transition into N_b is first order (even at infinite field) whereas the N_-/N_{+4} transition remains second order. It follows from our results that the N_{+4} region is broader in density and temperature at the moderate field $W/A_4 = 10$. Moreover, for $A_2 = 0.2A_4$ (when the parallel orientation of molecules is 4.4 times more energetically favorable than the perpendicular one) the N_{+4} phase exists at $W/A_4 = 5$; 10; 50, but is absent in infinite and zero fields. When the parallel orientation is almost 6 times more favorable than the perpendicular one ($A_2 = 0.3A_4$) the N_b phase finally excludes N_{+4}.

In conclusion, we have studied a model system of uniaxial particles with antagonistic molecular interactions in the presence of a disorienting field causing an easy-plane anisotropy. This may lead to spontaneous orientational ordering of two types. The first is a biaxial nematic N_b with a nonzero value of the conventional biaxial order parameter $S_x = \left\langle \sin^2\theta\cos(2\varphi)\right\rangle_\omega$ (as well as of the uniaxial order parameter $S_z = \langle P_2(\cos\theta)\rangle_\omega$). The biaxial order parameters is however zero in the second spontaneously ordered phase, named N_{+4}, which we predict to arise at intermediate density. The symmetry of this phase is defined instead by a nonvanishing higher orientational average $S_{xy} = \left\langle \sin^2\theta\cos(4\varphi)\right\rangle_\omega$. This corresponds to a cross-shaped distribution of preferred molecular orientations, which can be rotated in the xy-plane without any energy cost (just like the usual transverse director in the N_b phase). This feature is connected with Goldstone-mode fluctuations of particle orientations in the plane.

Device applications for nematic phases are of widespread importance [5], and a number have already been envisaged for the biaxial nematic phase [12]. But thermotropic biaxial nematics have proved to be very elusive. One of the causes is that they are displaced by other structures. It makes it necessary to study all phases nearby (like N_{+4}) and conditions of their existence. An important question that arises is how to look for the new N_{+4} phase experimentally. N_{+4} could be confused with N_b in its elastic behavior. Both should demonstrate elastic resistance to the following distortions: 1) splay and bend in the xy-plane, 2) twist in the z direction. In contrast to N_b and N_{+4}, the uniaxial phase N_- does not posess such elastic behavior. The new symmetry of the N_{+4} phase also should influence the behavior of defects, as is also true for biaxial nematics [13]. On the other hand, since it has a fourfold symmetry axis, the N_{+4} phase is uniaxial with respect to second rank tensors, and therefore not distinguishable by simple dielectric or optical measurements from an ordinary uniaxial nematic.

Finally it is worthwile to note that the N_{+4} symmetry could in principle appear without external fields, in molecules of a more complex geometry. From this point of view the investigation of systems of biaxial particles in search of the N_{+4} phase represents an interesting prospect. The ordering of long molecular axes eventually leads to the restriction of orientational phase space of the molecular short axes without any external field. This can influence their further orientational ordering with the same consequences as discussed above.

R. O. S. thanks NATO and the Royal Society for financial support.

---

[1] C. P. Fan and M. J. Stephen, Phys. Rev. Lett., **25**, 500 (1970)

[2] S. Varga, G. Cronome, and I. Szalai, Mol. Phys., **98**, 911 (2000); P. Palfy-Muhorray and D. A. Dunmur, Molec. Cryst. & Liq. Cryst. **97**, 337 (1983); T. G. Sokolovska, R. O. Sokolovskii, and M. F. Holovko, Phys. Rev. E **64**, 051710 (2001)

[3] Such effects might also be addressed within the Landau-de Gennes expansion of the free energy or some related approach. (See P. G. de Gennes and J. Prost *The Physics of Liquid Crystals*, 2nd Ed. (Clarendon, Oxford, 1993); and e.g. S. Hess, Z. Naturforsch **31a**, 1034 (1976). Such approaches do not exclude orientational interactions beyond P_2, but a microscopic calculation is needed to link these with the relevant expansion coefficients.

[4] J. J. Vroege and H. N. W. Lekkerkerker, Rep. Prog. Phys. **55**, 1241 (1992); R. Bruinsma, Phys. Rev. E **63**, 061705 (2001)

[5] R. Shashidhar (Ed), *Liquid Crystal Materials, Devices, and Displays* (Proc. SPIE, 1995)

[6] T. Swager and T. M. Long; Absts. Book, 19th Int. Liquid Crystal Conf., Edinburgh 2002, p. I20.

[7] C. G. Gray and K. E. Gubbins, *Theory of Molecular Fluids, vol.1: Fundamentals* (Clarendon, Oxford, 1984).

[8] J. Toner, Phys. Rev. A **27**, 1157 (1983); T. C. Lubensky and L. Radzihovsky, Phys. Rev. E **66**, 031704 (2002)

[9] T. G. Sokolovska and R. O. Sokolovskii, Phys. Rev. E **59**, R3819 (1999)

[10] J. A. C. Veerman and D. Frenkel, Phys. Rev A **45**, 5632 (1992)
[11] S. Romano, Liq. Cryst. **12**, 641 (1992)
[12] G. R. Luckhurst, Thin Solid Films **393**, 40 (2001)
[13] M. V. Kurik and O. D. Lavrentovich, Uspehi Phys. Nauk **154**, 381 (1988)